

CONCERNING THE NEGATIVE PART OF THE SPECTRUM OF
ONE-DIMENSIONAL AND MULTI-DIMENSIONAL
DIFFERENTIAL OPERATORS ON VECTOR-FUNCTIONS

I. M. Glazman


NASA-TT-F-14690) CONCERNING THE NEGATIVE
PART OF THE SPECTRUM OF ONE-DIMENSIONAL
AND MULTI-DIMENSIONAL DIFFERENTIAL
(Linguistic Systems, Inc., Cambridge,
Mass.) 10 p HC \$3.00 CSCL 12A

N73-26620

Unclass

G3/19 08708

Translation of: "Ob otritsatel'noy chasti
spektra odnomernykh i mnogomernykh differen-
tsial'nykh operatorov nad vektor-funktsiyami,"
Doklady Akademii Nauk SSSR, Vol. 119, Issue 3,
1958, pp. 421-424.



NATIONAL AERONAUTICS AND SPACE ADMINISTRATION
WASHINGTON, D.C. 20546

JULY 1973

CONCERNING THE NEGATIVE PART OF THE SPECTRUM OF
ONE-DIMENSIONAL AND MULTI-DIMENSIONAL
DIFFERENTIAL OPERATORS ON VECTOR-FUNCTIONS[†]

I. M. Glazman

The present note is devoted to an extension of theorems of [421*] note [1a], that supplements earlier obtained results [1b] concerning the spectrum of single-dimensional and multi-dimensional differential operators on vector-functions.

Let $\vec{\mathcal{L}}_2(0, \infty)$ be a Hilbert space of vector-functions $y(x) = \{y_k\}_{k=1}^m$ ($m < \infty$) with scalar product

$$(y, z) = \int_0^\infty \sum_{k=1}^m y_k(x) \overline{z_k(x)} dx,$$

and $l[y]$ be a differential operation of the form

$$l[y] = (-1)^n y^{(2n)} + Q(x)y \quad (0 \leq x < \infty), \quad (1)$$

where $Q(x)$ is a Hermite matrix-function of the m -th order. The least and, respectively, the greatest eigenvalue of matrix $Q(x)$ let us designate by $\mu(x)$ and $\nu(x)$. By \mathbb{L} let us designate any self-adjoint expansion of an operator with minimal area of determination generated in $\vec{\mathcal{L}}_2(0, \infty)$ by operation (1). The negative part of any function $f(x)$ let us designate by $f^*(x)$, such that $f^*(x) = \min\{0, f(x)\}$.

The use of lemma 1 of note [1a], where one must replace functional $\Phi_\varepsilon[y]$ by functional

[†]Presented by Academician S. N. Bernshteyn, October 24, 1957.

*Numbers in right hand margin indicate pagination of foreign text.

$$\Phi_\varepsilon[y] = \int_0^\infty \sum_{k=1}^m |y_k^{(n)}(x)|^2 dx + \int_0^\infty \sum_{j,k=1}^m Q_{jk}(x) y_j(x) \overline{y_k(x)} dx + \varepsilon \int_0^\infty \sum_{k=1}^m |y_k(x)|^2 dx,$$

leads to the following results.

Theorem 1. If in the case of any $\delta > 0$ is fulfilled the inequality

$$\int_{M_\delta} |\mu^*(x)| dx < \infty,$$

where M_δ is the set of values x , for which $|\mu^*(x)| \geq \delta$, then the negative part of the spectrum of operator \mathbb{L} is bounded below and discrete.

Assuming, further,

$$\alpha_n = \frac{(2n-1)!!}{2^n}, \quad A_n = (2n-1)^{-1} \left[\sum_{k=1}^n \frac{(-1)^{k-1} C_{n-1}^{k-1}}{2n-k} \right]^{-1} (n-1)!,$$

$$B_n^2 = \frac{n(4n^2-1)}{3 \cdot 4^{n-1}} \sum_{k=1}^n \frac{1}{2k-1} \sum_{h=0}^{2n-2} \frac{(-1)^h C_{2n-2}^h}{4n-3-k} \left[\sum_{h=1}^n \frac{(-1)^{h-1} C_{n-1}^{h-1}}{2n-k} \right]^{-2},$$

let us mention the following two theorems.

Theorem 2. The negative part of the spectrum of operator \mathbb{L} consists of a finite number of eigenvalues, if one of the following conditions is fulfilled:

1. $\mu(x) \geq -\alpha_n^2 x^{-2n}$ in the case of large x .
2. In the case of any $\delta > 0$

$$\int_{M_\delta} x^{2n-1} |\mu^*(x)| dx < \infty,$$

where M_δ is the set of values x for which

$$|\mu^*(x)| \geq (\alpha_n^2 - \delta) x^{-2n}.$$

3. In the case of some $p \geq 1$

$$\int_0^{\infty} x^{2np-1} |\mu^*(x)|^p dx < \infty.$$

Theorem 3. The negative part of the spectrum of operator L is an infinite set if one of the following conditions is fulfilled:

1. In case of some $\delta > 0$ and large x

$$v(x) < -(\alpha_n^2 + \delta) x^{-2n}.$$

2. $v(x) \leq 0$ in case of large values of x and

$$\liminf_{\rho \rightarrow \infty} \rho^{2n-1} \int_{\rho}^{\infty} |v(x)| dx > A_n^2.$$

3. $v(x) \leq -\alpha_n^2 x^{-2n}$ in the case of large x and

$$\liminf_{\rho \rightarrow \infty} \ln \rho \int_{\rho}^{\infty} x^{2n-1} |v(x) + \alpha_n^2 x^{-2n}| dx > B_n^2.$$

4.

$$\int_0^{\infty} v(x) dx = -\infty.$$

In conditions 2 and 3 one can replace $\liminf_{\rho \rightarrow \infty}$ by $\lim_{\rho_k \rightarrow \infty}$.

Theorems 1-3 are connected with the oscillation properties of the system of differential equations

$$(-1)^n y^{(2n)} + Q(x)y = \lambda y \quad (\lambda \leq 0),$$

that were studied by Sternberg [2] in the case of $n = 1$. Conditions 1 of theorem 2 and 1 of theorem 3 give an extension of the well-known theorem of Knezer concerning the oscillation of solutions of a differential equation of the second order. In the case of $n = 1$ one can obtain the following refinement of condition 1 of theorem 3 that for the case of a differential equation of the

second order was given by Hille [3] (see also [4]).

Theorem 4. If in the case of some $\delta > 0$ and some natural r for all sufficiently large values of x occurs the inequality

$$v(x) < -\frac{1}{4x^2} - \frac{1}{4x^2 \ln^2 x} - \dots - \frac{1+\delta}{4x^2 \ln^2 x \dots \ln^r x},$$

where $\ln_k x = \ln \ln_{k-1} x$, then the negative part of the spectrum of operator L consists of a finite number of eigenvalues.

The presented results are partially extended to multi-dimensional differential operations on vector-functions of the form

$$L[u] = -\Delta u + Q(P)u, \quad (2)$$

where P is a point of n -dimensional Euclidean space E_n ; $Q(P)$ is a Hermitian matrix-function of the m -th order determined in all E_n .

Operation (2) generates in Hilbert space $L_2(E_n)$ of vector-functions $u(P) = \{u_k(P)\}_{k=1}^m$ with scalar product

$$(u, v) = \int_{E_n} \sum_{k=1}^m u_k(P) \overline{v_k(P)} d\omega_P,$$

some differential operator L with a minimum area of determination.

Let $\mu(P)$ be the least eigenvalue of matrix $Q(P)$ and $\mu^*(P) = \min\{0, \mu(P)\}$. Let us present, for example, a formulation of the theorem that corresponds to theorem 1, and let us prove it for $m = 1$, $n = 2$ (in this case $\mu(P) = Q(P)$).

Theorem 5. If in the case of $\delta > 0$ the integral

$$\int_0^\infty |\mu_\delta^*(P)| dr,$$

where

$$\mu_\delta^*(P) = \begin{cases} \mu^*(P), & |\mu^*(P)| > \delta, \\ 0, & |\mu^*(P)| \leq \delta, \end{cases}$$

converges uniformly along angular coordinates, then:

1) operator L with a minimum area of determination (see [1c]) is self-adjoint;

2) the negative part of the spectrum of operator L is semi-bounded below and discrete (that is, consists of eigenvalues of finite multiplicity with a single possible limiting point $\lambda = 0$).

Proof. Converting the quadratic functional

$$\Phi_\varepsilon[u] = \iint_{(G)} |\nabla u|^2 r dr d\varphi + \iint_{(G)} Q|u|^2 r v dr d\varphi + \varepsilon \iint_{(G)} |u|^2 r dr d\varphi$$

to any finite function $u \in D_L$ with the help of the substitution of variables $u/\sqrt{r} = v$, let us obtain

$$\Phi_\varepsilon[u] = \iint_{(G)} |\nabla v|^2 dr d\varphi + \iint_{(G)} \left[Q(r, \varphi) + \frac{1}{4r^2} \right] |v|^2 dr d\varphi + \varepsilon \iint_{(G)} |v|^2 dr d\varphi.$$

In the case of randomly assigned ε ($0 < \varepsilon < 1$) let us select a number N such that there would be

$$\int_N^\infty |Q^*(r, \varphi)| dr < \frac{\varepsilon}{4},$$

and let us show that the functional

$$\Phi_\varepsilon[u] = \int_0^{2\pi} d\varphi \int_N^\infty \left\{ |\nabla v|^2 + \left[Q^*(r, \varphi) + \frac{1}{4r^2} + \varepsilon \right] |v|^2 \right\} dr$$

is nonnegative on any finite function $v \in D_L$ that equals zero in the range $r < N$.

From the Cauchy-Bunyakovsky inequality it follows that

$$\int_0^{2\pi} \int_N^\infty |\nabla v|^2 dr d\varphi \geq \int_0^{2\pi} d\varphi \int_N^\infty \left| \frac{\partial v}{\partial r} \right|^2 dr \geq \frac{1}{4} \left\{ \int_0^{2\pi} |\hat{v}(\varphi)|^2 d\varphi \right\}^2,$$

where the function $v(r, \phi)$ is normalized by the condition

$$\int_0^{2\pi} d\varphi \int_N^\infty |v|^2 dr = 1$$

and

$$\hat{v}(\varphi) = \max_{N \leq r < \infty} |v(r, \varphi)|.$$

Let us consider two cases separately:

$$1. \int_0^{2\pi} |\hat{v}(\varphi)|^2 d\varphi \leq \varepsilon. \quad 2. \int_0^{2\pi} |\hat{v}(\varphi)|^2 d\varphi > \varepsilon.$$

In the first case

$$\begin{aligned} \Phi_\varepsilon[u] &\geq \int_0^{2\pi} \int_N^\infty Q^*(r, \varphi) |v|^2 dr d\varphi + \varepsilon \int_0^{2\pi} \int_N^\infty |v|^2 dr d\varphi \geq \\ &\geq - \int_0^{2\pi} |\hat{v}(\varphi)|^2 \int_N^\infty Q^*(r, \varphi) dr d\varphi + \varepsilon \geq \varepsilon \left(1 - \frac{\varepsilon}{4}\right) > 0. \end{aligned}$$

In the second case

$$\Phi_\varepsilon[u] \geq \frac{1}{4} \left\{ \int_0^{2\pi} |\hat{v}(\varphi)|^2 d\varphi \right\}^2 + \int_0^{2\pi} \int_N^\infty Q^*(r, \varphi) |\hat{v}(\varphi)|^2 dr d\varphi,$$

such that

$$\Phi_\varepsilon[u] \geq \frac{1}{4} \int_0^{2\pi} |\hat{v}(\varphi)|^2 d\varphi \left[\int_0^{2\pi} |\hat{v}(\varphi)|^2 d\varphi - \varepsilon \right] > 0,$$

and inequality $\Phi_\varepsilon[u] \geq 0$ is established.

From the given inequality first of all follows [1d] the semi-boundedness of operator L below, and from that, according to a theorem of A. Ya. Povzner, the self-adjointness of operator L follows.

Further from this inequality on the basis of [1d] and of lemma 1 [1a] let us conclude that the negative part of the spectrum of operator L is discrete. The theorem is proven.

V. I. Lenin Kharkov Polytechnical Institute
Received October 24, 1957

REFERENCES

1. I. M. Glasman, a) DAN, Vol. 118, No. 3 (1958); b) Uch. zap. Khar'kovsk. gos. ped. inst., Vol. 18, mathematical series (1956); c) Tr. Khar'kovsk. nolitexhn. inst., Vol. 5, series engineering-physics, in 1 (1955); d) Matem. sborn., Vol. 35(77), 2, (1954).
2. R. Sternberg, Duke Math. Journal, Vol. 19, 311 (1952).
3. E. Hille, Trans. Am. Math. Soc., Vol. 64, No. 234 (1948).
4. R. Bellman, Theory of Stability [Teoriya ustoychivosti], Foreign Literature Press, 1954.
5. A. Ya. Povzner, Mathematical Collection [Matem. sborn.], Vol. 32(74), 1 (1953).

1. Report No. NASA TT F-14,690	2. Government Accession No.	3. Recipient's Catalog No.	
4. Title and Subtitle CONCERNING THE NEGATIVE PART OF THE SPECTRUM OF ONE-DIMENSIONAL AND MULTI- DIMENSIONAL DIFFERENTIAL OPERATORS OVER VECTOR-FUNCTIONS		5. Report Date July 1973	
7. Author(s) I. M. Glazman		6. Performing Organization Code	
9. Performing Organization Name and Address LINGUISTIC SYSTEMS, INC. 116 AUSTIN STREET CAMBRIDGE, MASSACHUSETTS 02139		8. Performing Organization Report No.	
12. Sponsoring Agency Name and Address NATIONAL AERONAUTICS AND SPACE ADMINISTRATION WASHINGTON, D. C. 20546		10. Work Unit No.	
		11. Contract or Grant No. NASW-2482	
		13. Type of Report & Period Covered TRANSLATION	
		14. Sponsoring Agency Code	
15. Supplementary Notes Translation of: "Ob otritsatel'noy chasti spektra odnomernykh i mnogomernykh differentsial'nykh operatorov nad vektor-funktsiyami," Doklady Akademii Nauk SSSR, Vol. 119, Issue 3, 1958, pp. 421.424.			
16. Abstract The article extends theorems of previous work by I.M. Glasman concerning the spectrum of one-dimensional and multi-dimensional differential operators on vector functions. Given a Hilbert space of vector functions and a differential operator of specific form, it is first shown that the negative part of the spectrum of any self-adjoint expansion of an operator L is discrete and bounded below (Theorem 1). Further specification of two separate sets of conditions causes the negative part of the spectrum of operator L (1) to consist of a finite number of eigenvalues (Th. 2) and (2) to be an infinite set (Th. 3). These 3 theorems are connected with the oscillation properties of certain differential equations studied by Sternberg [2]. Refinement of one condition of Theorem 3 applied to a differential equation of second order results in the negative part of the spectrum of operator L consisting of a finite number of eigenvalues (Th. 4). Then it is proven that for specific integral types operator L with a minimum area of determination is self-adjoint and (cont)			
17. Key Words (Selected by Author(s))		18. Distribution Statement UNCLASSIFIED - UNLIMITED	
19. Security Classif. (of this report) UNCLASSIFIED	20. Security Classif. (of this page) UNCLASSIFIED	21. No. of Pages 10	22. Price

the negative part of the spectrum of operator L is discrete and semi-bounded below.